



[ANIMAL HEALTH & WELFARE \(/ADVOCATE/CATEGORY/ANIMAL-HEALTH-WELFARE\)](#)

---

# Evaluating heritability of growth, cold tolerance in Chinese white shrimp

Monday, 15 July 2019

By Dr. Xianhong Meng

**Results showed low heritability, low correlation between these traits**



This study estimating genetic heritabilities for body weight and length for Chinese white shrimp showed low heritability for both traits and a low correlation between growth and cold tolerance traits, Photo by Darryl Jory.

*Editor note: This article has nine co-authors (listed in the tags below), but we only include affiliation information for the corresponding author, Dr. Xianhong Meng. For more information, refer to [original publication \(https://doi.org/10.1371/journal.pone.0183801\)](https://doi.org/10.1371/journal.pone.0183801).*

The Chinese white shrimp (*Fenneropenaeus chinensis*) – also known as oriental or fleshy prawn – is commercially cultivated in China. The species is one of the most representative indigenous aquaculture species in the country, it is mainly distributed in the Yellow Sea and Bohai Sea area, and cultured in Shandong, Hebei, Liaoning, Tianjin, Jiangsu offshore, as well as the west and south coasts of the Korean Peninsula. Its optimum water temperatures range from 18 to 30 degrees-C – although it tolerates 4 to 38 degrees-C – with an optimum temperature of around 25 degrees-C.

Hatchery production of seedstock started developing in the 1970s, and the culture area expanded through the 1980s. From 1988 to 1993, annual farmed production of *F. chinensis* yielded as much as over 200,000 metric tons (MT), with very significant economic benefits to the aquaculture industry. However, in 1993 the *F. chinensis* farming industry has hit by the white spot syndrome virus (WSSV) and suffered serious economic losses, with 1994 national production falling sharply to around 60,000 MT. This led to significant research efforts that resulted in the development of three new lines or varieties of Chinese white shrimp, namely the Huanghai No. 1 in 2003, the Huanghai No. 2 in 2008, and the Huanghai No. 3 in 2013, which enabled a gradual recovery of China's farming of *F. chinensis*.

However, in recent years, the shrimp farming industry is facing new problems. The intensifying, abnormal climate – especially repeated cold spells nationwide – has resulted in significant shrimp mortality and serious economic losses to shrimp farmers. It is expected that the production of more cold-tolerant shrimp species like *F. chinensis* will help alleviate this issue.

This article – adapted and summarized from the **original publication** (<https://doi.org/10.1371/journal.pone.0183801>) – presents results of an evaluation of the growth and cold tolerance traits of *F. chinensis* juveniles from 99 different families. The study used an indoor artificial cold tolerance challenge with analysis of heritability of body weight (BW), body length (BL) and other parameters. This evaluation would provide a theoretical basis for breeding more cold tolerance varieties of *F. chinensis*.

## Study setup

The experiment was conducted at the Mariculture Research Station of the Yellow Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences in Qingdao City, Shandong Province, China. The experimental materials were 99 Huanghai No. 2 families of *F. chinensis* developed in 2015 (including 12 half-sib families), representing the G10-generation, at age of 40 days.

For the cold tolerance challenge treatment, 30 juvenile shrimp (average body weight of  $0.073 \pm 0.043$  grams and body length of  $19.790 \pm 3.607$  mm, respectively) were randomly sampled from each family and placed in water in aerated storage boxes (26.5 x 20 x 16.5 cm). The boxes were placed in a refrigerator adjustable temperature. The refrigerator temperature was initially dropped to 14 degrees-C at a rate of 2 degrees-C daily, and then further dropped to (and maintained at) 4 degrees-C at a rate of 2 degrees-C daily, until all the juvenile shrimps died. The juvenile shrimp were fed four times daily and 30 percent of the water exchanged every day. Dead shrimps were collected every two hours and information data collected on family, temperature at mortality, and body weight and body length.

For detailed information on the experimental design and setup; the cold tolerance challenge treatment of the juveniles, data processing; estimates of heritability and correlation analysis of growth traits; and correlation analysis of growth traits and cold tolerance traits in *F. chinensis* juveniles, please consult the original publication.

## Results and discussion

The sample size, mean, maximum, minimum, standard deviation and coefficient variation for growth and cold tolerance traits of *F. chinensis* are shown in Table 1. The mean body weight (BW), body length (BL), temperature at death (TAD), cooling degree hours (CDH) and survival rate for each family at half lethal time (SR<sub>50</sub>) were 0.073 grams, 19.790 mm, 6.506 degrees-C, 333.937 degrees-C\*h, and 48.950 percent, respectively, with coefficients of variation ranging from 18.226 percent to 59.919 percent.

## Wang, cold tolerance, Table 1

Traits	Number	Mean	Minimum	Maximum	Coefficient of variation (%)
Body weight (g)	2,508	0.073	0.005	0.370	59.919
Body length (mm)	2,508	19.79	9.610	33.120	18.226
Cooling degree hours (degrees-C*hour)	2,508	333.937	0.800	720.000	48.221
Temperature at death (degrees-C)	2,508	6.506	3.000	14.000	36.241
Half lethal time, SR50 (%)	99	48.95	0.000	100.000	54.107

Table 1. Description statistics of growth and cold tolerance traits in *F. chinensis* juveniles. Adapted from original publication.

The coefficient of variation of body weight was the greatest (59.919 percent), indicating that there was great variance in body weight between the different families. While the coefficient of variation of body length was the smallest (18.226 percent), indicating that there was comparatively small variance in body length between different families. Among all the cold tolerance traits, the SR<sub>50</sub> had the highest phenotypic variance, with 54.107 percent coefficient variation. It was followed by CDH (48.221 percent), and by TAD with the lowest (36.241 percent).

In the cold tolerance challenge experiment, mortality of shrimp was observed when water temperature was 13.9 degrees-C and the CDH was 0.800 degrees-C\*h. All the shrimps had died by the time the CDH reached 720 degrees-C\*h (Fig 1). The cumulative mortality curve is shown in Fig 1.




Fig 1. Cumulative mortality of *F. chinensis* juveniles at different CDH values during the cold tolerance challenge.

Regarding the heritability for growth traits of *F. chinensis* juveniles, our research results showed that the heritabilities for body weight and body length were low ( $0.078 \pm 0.124$  and  $0.131 \pm 0.133$ , respectively). The heritabilities for growth traits obtained in various studies by other researchers were higher than those obtained in our study because the heritability estimates were affected by different shrimp varieties or different growth periods (age) of the same variety, and by the different genetic backgrounds and structures of the breeding populations.

TAD and CDH were the parameters used to estimate heritability for cold tolerance in our study. The heritability estimates for TAD and CDH were  $0.265 \pm 0.091$  and  $0.077 \pm 0.058$ , respectively, which were moderate and low in magnitude. The heritability for cold tolerance traits of *F. chinensis* juveniles have not been previously reported, but Li Wenjia and co-workers have previously evaluated the heritability of cold tolerance traits like CDH in Pacific white shrimp (*Litopenaeus vannamei*) adults and reported it as low with an estimate of  $0.0258 \pm 0.0205$ .

Regarding the correlation analysis of growth and cold tolerance traits of *F. chinensis* juveniles, our results showed that the phenotypic correlations between growth and cold tolerance were low. The genetic correlation has a larger range ( $0.0526 \sim 0.9914$ ), and all were not significantly different from zero due to large standard errors. The results essentially agreed with the results of other researchers, who studied genetic correlation between growth and cold tolerance traits of *L. vannamei* and genetic correlation between body weight and cold tolerance traits of Nile tilapia, respectively.

## Perspectives

Results from this study showed that there was a low correlation between growth and cold tolerance traits, indicating that growth and cold tolerance should be considered collectively in the breeding program of shrimp.

Further studies are required for the correlation of different traits of *F. chinensis*, so as to provide more accurate data for breeding of cold tolerance *F. chinensis*. In addition, based on traditional breeding methods, traits of low heritability could be improved by means of studies of molecular biology to accelerate the breeding process.

## Author

---



### DR. XIANHONG MENG

Corresponding author

Key Laboratory of Sustainable Utilization of Marine Fisheries Resources, Ministry of Agriculture, Yellow Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Qingdao, China; and Laboratory for Marine Fisheries Science and Food Production Processes, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China; and Shanghai Ocean University, Shanghai, China  
[mengxianhong@ysfri.ac.cn](mailto:mengxianhong@ysfri.ac.cn) (mailto:mengxianhong@ysfri.ac.cn).

7/16/2019

Evaluating heritability of growth, cold tolerance in Chinese white shrimp « Global Aquaculture Advocate

Global Aquaculture Alliance