



[FEED SUSTAINABILITY \(/ADVOCATE/CATEGORY/FEED-SUSTAINABILITY\)](#)

Feed study examines effects of fishmeal storage

Thursday, 1 November 2012

By Dong-Fang Deng, Ph.D. , Zhi Yong Ju, Ph.D. , Lytha D. Conquest , Peter J. Bechtel, Ph.D. and Scott Smiley, Ph.D.

Storage conditions change fishmeal composition, not shrimp growth

Fishmeal is an important dietary ingredient for aquatic diets due to its ideal amino acid profile for most cultured species, excellent palatability and the availability of polyunsaturated fatty acids and minerals. The quality of fishmeal, however, can be affected by different factors.

Many studies have shown the quality of fishmeal is affected by raw material freshness as well as the species of fish and processing conditions used to manufacture the fishmeal. Different responses to the quality of fishmeal have also been attributed to the species being tested and their size or life history stage.

Furthermore, management of the finished meals is critical for the quality of fishmeal, which is susceptible to oxidation. Both rancidity from lipid oxidation and the presence of biogenic amines produced from protein oxidation affect fishmeal quality.

Responses and sensitivity to the effects of oxidized fishmeal are different for different species of fish and shrimp. For example, cadaverine and putrescine act as chemoattractants for freshwater prawns at 2,000 mg/kg in diets. Putrescine was found to adversely affect both feed intake and growth of rainbow trout at a concentration of 13,300 mg/kg diet. Therefore, evaluation using a growth trial is necessary to evaluate the quality of a fishmeal in addition to data representing the degree of oxidation based on chemical analysis.



Juvenile white shrimp received feeds containing fishmeal stored for varied time periods under different temperatures.

Pink salmon

Pink salmon is one of the major harvested species in commercial fisheries in Alaska, USA. The total harvest for 2011 was about 115 million fish weighing about 180,000 metric tons. The processing of this fish generates significant waste reflecting 25 to 30 percent of the whole fish, which is available for the production of fishmeal. A 2009 study by Ted Wu and fellow researchers found that delayed handling of raw pink salmon processing by-products impaired the quality of fishmeal based on chemical analysis related to lipid and protein oxidation.

In the authors' study, the objective was to investigate the effects of storage temperature and duration on the quality of finished fishmeal based on the growth performance of juvenile shrimp. The study was funded by a grant from the U.S. Department of Agriculture Agricultural Research Service and through a cooperative agreement with the University of Alaska Fairbanks.

Growth trial

Pink salmon meal from the Kodiak Fishmeal Co. in Alaska was processed by the standard Atlas-Stord wet reduction technology. The fishmeal was divided into equal proportions and sealed in different bags protected from light under the conditions shown in Table 1.

Fang-Deng, Conditions for fishmeal, Table 1

FM6-CT	FM12-CT	FM6-AT	FM12-AT	FM-C
6 months @ 4° C	12 months @ 4° C	6 months @ 30° C	12 months @ 30° C	12 months @ -80° C

Table 1. Conditions for fishmeal storage.

Six test diets were formulated as 36.8 percent whole wheat, 25.0 percent soybean meal, 15.0 percent fishmeal, 6.0 percent squid meal and 17.8 percent other products, including minerals, vitamins and oil. All tests diets had similar 32 percent protein and 9 percent lipid contents. The diets were agglomerated by a pellet mill into 2.4- x 4.0-mm pellets.

The growth trial was conducted in an indoor flow-through saltwater system with a photoperiod of 12 hours light and 12 hours dark. Specific pathogen-free shrimp were randomly stocked into 20 tanks at a density of 20 shrimp/tank (83 shrimp/m²). The shrimp were fed daily at 8 to 15 percent of body weight using automatic feeders. The feed for each tank was loaded into the feeder at 8:30 a.m. daily. Shrimp were weighed every two weeks, and the feeding rate was adjusted based on growth rate.

During the trial, water quality was monitored, with temperature at 26.5 ± 0.1 degrees-C; dissolved-oxygen concentrations at 6.6 ± 0.2 mg/L, salinity at 30.7 ± 0.1 ppt and pH at 7.6 ± 0.2 . Total ammonia-nitrogen levels were below 0.05 ± 0.01 mg/L. At the end of a six-week feeding trial, shrimp from each tank were killed with ice. The tail muscles from shrimp in each tank were pooled, freeze dried and then stored at -80 degrees C until analyzed.

All data were subjected to one-way analysis of variance. Treatment means were considered significantly different at the level of $P < 0.05$.

Results

The proximate compositions of the salmon meals stored at different conditions were similar except that the two meals stored at 4 degrees-C showed increased moisture levels and thus had lower levels of dry matter (Table 2). This was likely due to absorption of moisture in the fishmeal during the storage period.

Fang-Deng, Proximate and amino acid compositions, Table 2

Analysis	FM6-CT	FM12-CT	FM6-AT	FM12-AT	FM-C
Proximate Composition (%)					
Ash	87.60	87.00	96.30	96.20	91.90
Dry matter	12.90	11.90	13.80	13.10	12.60
Crude protein	59.50	60.80	61.70	61.30	60.60
Crude lipid	9.40	9.40	9.80	9.10	9.50
Non-Essential Amino Acids (%)					
Alanine	4.65	4.51	4.50	4.42	4.69
Aspartate + asparagine	3.84	3.62	3.82	3.53	4.30
Glutamate + glutamine	4.42	4.66	5.03	5.07	5.15
Glycine	7.26	7.24	6.75	6.54	7.35
Proline	3.05	3.25	3.06	3.06	3.15
Serine	2.19	2.21	2.08	2.09	2.36
Tyrosine	1.83	1.69	1.47	1.58	1.56
Essential Amino Acids (%)					
Arginine	5.37	5.26	5.24	5.16	5.56
Histidine	2.68	2.48	2.61	2.61	2.71
Isoleucine	2.72	2.70	2.49	2.50	2.71
Leucine	3.14	3.12	2.69	2.95	3.15
Lysine	3.69	3.70	2.98	3.01	3.71
Methionine	1.45	1.44	1.46	1.34	1.49
Phenylalanine	1.87	1.88	1.25	1.12	1.80
Threonine	2.75	2.59	2.71	2.62	2.82
Valine	4.75	4.29	3.87	3.99	4.23
Subtotal of Non-Essential Amino Acids (%)	27.24	27.18	26.73	26.30	28.56
Subtotal of Essential Amino Acids (%)	28.42	27.45	25.30	25.30	28.17
Total	55.66	54.64	52.03	51.60	56.73

Table 2. Proximate and amino acid compositions of fishmeals used in test diets.

The overall levels of amino acids were lower for the salmon meal stored at 30 degrees-C than those of fishmeal preserved at 4 or minus-80 degrees-C. This indicated degradation or oxidation of amino acids in the fishmeal stored under high temperature. The proximate compositions were very similar among different test diets (Table 3).

Fang-Deng, Nutrient compositions, Table 3

Nutrient	FM6-CT	FM12-CT	FM6-AT	FM12-AT	FM-C
Dry matter	92.9	93.4	93.3	93.5	93.4
Ash	12.9	13.0	13.0	12.9	23.0
Protein	31.6	31.8	32.3	31.1	31.4
Lipid	9.1	8.8	8.9	9.0	9.3
Gross energy (kcal/100 g)	410.6	412.4	412.6	413.9	410.9

Table 3. Nutrient compositions (%) of test diets.

Results from the growth trial showed that shrimp fed diets formulated with different fishmeal had similar growth rates, feed-conversion ratios and survival (Table 4). The overall survival of shrimp was similar among different treatments, but at 75 to 85 percent was lower than the 90 percent survival observed in previous studies by the authors using the same reference diet.

Fang-Deng, Mean growth performance, Table 4

Parameter	FM6-CT	FM12-CT	FM6-AT	FM12-AT	FM-C
Initial body weight (g)	1.17 ± 0.02	1.16 ± 0.01	1.13 ± 0.03	1.17 ± 0.01	1.17 ± 0.03
Final body weight (g)	6.80 ± 0.20	6.60 ± 0.10	6.50 ± 0.10	7.20 ± 0.30	6.90 ± 0.20
Weight gain/week (g)	0.94 ± 0.03	0.91 ± 0.02	0.90 ± 0.02	1.01 ± 0.05	0.95 ± 0.03
Specific growth rate (%)	4.19 ± 0.05	4.15 ± 0.06	4.17 ± 0.03	4.33 ± 0.11	4.22 ± 0.05
Feed-conversion ratio	2.08 ± 0.03	2.10 ± 0.05	2.08 ± 0.04	2.01 ± 0.07	2.05 ± 0.03
Survival (%)	81.30 ± 3.20	85.00 ± 6.10	82.50 ± 4.30	75.00 ± 7.10	85.00 ± 2.00
Dry matter (%)	23.30 ± 0.20	23.10 ± 0.30	22.80 ± 0.20	23.10 ± 0.30	22.90 ± 0.20
Ash (%)	1.45 ± 0.05	1.30 ± 0.06	1.32 ± 0.03	1.36 ± 0.03	1.31 ± 0.03
Protein (%)	20.30 ± 0.10	19.90 ± 0.30	19.40 ± 0.40	20.00 ± 0.40	19.80 ± 0.70
Lipid (%)	0.98 ± 0.02	0.97 ± 0.04	0.92 ± 0.05	0.96 ± 0.02	0.93 ± 0.05
Gross energy (cal/g)	114.60 ± 1.10	111.90 ± 2.10	108.50 ± 0.90	113.30 ± 2.60	109.90 ± 2.40

Table 4. Mean growth performance of shrimp and nutritional composition of tail muscle of shrimp fed the test diets.

The reason may be the higher stocking density used in this trial. The quality of shrimp from a different family used in different studies may have also contributed to the different observation. The nutritional composition of shrimp tail muscle was similar among the different dietary treatments.

Perspectives

Results of this study suggested that salmon meal stored for up to 12 months at 30 degrees-C tended to show protein degradation. The study did not find any significant effect of the different types of fishmeal on growth performance, feed utilization or the nutritional composition of shrimp muscle. Therefore, the quality of shrimp feed was not downgraded when 15 percent of the fishmeal was included in a diet under current testing conditions.

However, the observation might have been different if a higher level of the aged fishmeal was tested in the diet and also if it was tested with shrimp of different sizes. A further study with a longer feeding trial will be needed for a more complete evaluation of the nutritional quality of aged fishmeal.

(Editor's Note: This article was originally published in the November/December 2012 print edition of the Global Aquaculture Advocate.)

Authors



DONG-FANG DENG, PH.D.

Aquatic Feeds and Nutrition Department

Oceanic Institute

41-202 Kalaniana'ole Highway

Waimanalo, Hawaii 96795 USA

dfdeng@oceanicinstitute.org (<mailto:dfdeng@oceanicinstitute.org>)



ZHI YONG JU, PH.D.

Aquatic Feeds and Nutrition Department
Oceanic Institute
41-202 Kalaniana'ole Highway
Waimanalo, Hawaii 96795 USA



LYTHA D. CONQUEST

Warren G. Dominy, Ph.D.
Aquatic Feeds and Nutrition Department
Oceanic Institute
41-202 Kalaniana'ole Highway
Waimanalo, Hawaii 96795 USA



PETER J. BECHTEL, PH.D.

United States Department of Agriculture
Agricultural Research Service
Subarctic Agricultural Research Unit
Fishery Industrial Technology Center
Kodiak, Alaska, USA



SCOTT SMILEY, PH.D.

Fishery Industrial Technology Center
School of Fisheries & Ocean Sciences
University of Alaska
Kodiak, Alaska, USA

Copyright © 2016–2019
Global Aquaculture Alliance