



**Alliance**

(<https://www.aquaculturealliance.org>).

---



**Global  
Aquaculture  
Advocate**<sup>™</sup>

Intelligence

---

# Pacific oysters, abalone undergo sensory evaluation

Monday, 1 November 2010

By Nick Elliott , Conor Delahunty , Ciarán Forde , Maeva Cochet and Peter Kube

**Results demonstrate ability to differentiate among regions, farms and growing conditions**



Sensory evaluation is a scientific approach to measurement of how food products stimulate our senses.

Sensory evaluation is a scientific approach to the measurement and analysis of how food and beverage products stimulate our senses. It is used routinely to guide food product research and development, production, manufacture and handling, and quality control.

Information on the sensory properties of food and beverages can be related to differences in their chemical and physical compositions. The sensory differences also can be related to differences in consumer preference, offering insight on what drives product acceptance.

## Seafood evaluation

A pilot study on the sensory properties of Pacific oysters and abalone was conducted at the CSIRO Sensory Laboratory in Sydney, Australia, by a team of scientists from CSIRO Food and Nutritional Sciences. It found differences among the sensory properties of samples produced in different growing conditions and regions.

The pilot study began by establishing protocols for the preparation and tasting of the oysters and abalone, as well as a sensory evaluation vocabulary for odor, texture, flavor and aftertaste for each product. Once consensus was reached, the samples underwent an objective sensory assessment by a panel of 10 trained assessors, whose role was to quantitatively rate the perceptual differences among the samples.

The preparation protocols set guidelines for thawing, shucking, washing, steaming for abalone and plating. The tasting protocols ensured panel members applied a consistent approach to sniffing, biting and chewing the food samples during the evaluation.

To find meaningful perceptual differences, the samples were sourced from three regions of Australia. They were not meant to represent a robust regional comparison. The samples of Pacific oysters were sourced from farms in New South Wales, Tasmania and South Australia. The abalone came from farms in South Australia, Tasmania and Victoria, and wild harvest in Tasmania.



Differences related to region and growing conditions could be identified in the tests.

## Differences perceived

The evaluation results demonstrated the ability of sensory testing to differentiate among regions, farms and growing conditions for both oysters and abalone.

All 12 of the attributes measured in oysters showed significant differences related to growing regions (Fig. 1). In addition, the abalone samples differed according to growing and harvesting conditions. The larger, wild-harvested abalone were perceived as having greater chewiness, firmness and springiness than farmed abalone, and containing higher levels of earthy and metallic flavors.




Fig. 1: Perceived intensity differences of key sensory attributes among Pacific oysters from three farms.

Farming systems and diets also influenced the sensory properties of farmed abalone samples, as shown in Figure 2. For example, abalone from one farm was called significantly more savory and salty. The panel was also able to significantly differentiate abalone of the same cohort but on different diets at one farm.



## Authors

---



### **NICK ELLIOTT**

CSIRO Food Futures Flagship  
CSIRO Marine and Atmospheric Research  
GPO Box 1538, Hobart  
Tasmania 7001 Australia

[nick.elliott@csiro.au](mailto:nick.elliott@csiro.au) (<mailto:nick.elliott@csiro.au>).



### **CONOR DELAHUNTY**

CSIRO Food Futures Flagship  
CSIRO Food and Nutritional Sciences  
North Ryde, New South Wales, Australia



### **CIARÁN FORDE**

CSIRO Food Futures Flagship  
CSIRO Food and Nutritional Sciences  
North Ryde, New South Wales, Australia



### **MAEVA COCHET**

CSIRO Food Futures Flagship  
CSIRO Food and Nutritional Sciences  
North Ryde, New South Wales, Australia



**PETER KUBE**

CSIRO Food Futures Flagship  
CSIRO Marine and Atmospheric Research  
Hobart, Tasmania, Australia

Copyright © 2016–2019 Global Aquaculture Alliance

All rights reserved.