



Alliance

(<https://www.aquaculturealliance.org>).




Health & Welfare

Parasite management important to future cod aquaculture in Scotland

Thursday, 1 December 2005

By Rachel Kilburn , Campbell Pert , Dr. David Bruno and Dr. Rob Raynard

Site-specific health management, biosecurity protocols essential



Once settled on the gills of a cod, the parasitic copepod *L. branchialis* can also damage the heart of the fish.

Atlantic cod (*Gadus morhua*), one of the leading food fish in the United Kingdom, have traditionally been fished in Scottish waters. Currently 270,000 metric tons (MT) of *G. morhua* are consumed every year in the U.K., but with decreasing wild stocks, this demand could in part be met through the cultivation of the species.

Wild Atlantic cod have an exceptionally varied parasite fauna, most of which causes little or no effect on the fecundity and health of the fish. The distribution of parasites within wild populations of cod are not random, but tend to be dispersed. The ecological processes that determine parasite distribution in wild cod are likely to result in different distribution and abundance of the parasites in farmed cod.

Densely farmed cod may be at greater risk if environmental conditions favor certain parasites and allow their populations to increase to a high level. However, farmed cod may have reduced loads of parasites if the cod can be isolated from intermediate hosts that carry infective stages of the parasites.

It is difficult to predict which parasites of the many naturally occurring species may arise as a problem to future cod aquaculture. Therefore, parasitologists at the FRS Marine Laboratory in Aberdeen, Scotland, are conducting baseline prevalence studies of parasites in wild cod and identifying those that may trouble cod cultivation in the future.

Transmission



Black spot is a common parasite that manifests itself on the skin of species like cod and this whiting.

Principal factors in the transmission of parasites between wild and farmed cod will be whether the parasites have direct monoxenous life cycles or indirect heteroxenous life cycles with many intermediate hosts, and the distribution of these hosts within the vicinity of fish farms.

Many cod parasites possess complex life cycles with stages that infect definitive hosts where maturation and sexual reproduction take place or involve asexual proliferation in intermediate hosts such as aquatic snails. Transmission between definitive and intermediate hosts can occur through free-living infective stages, as seen with sea lice, *Caligus elongates*; or predation by one host on the previous host in the life cycle, as with *Anisakis* sp. and *Cryptocotyle lingue* (black spot).

Table 1. Life cycles of cod parasites.

Species	Life Cycle	Site of Infection
Nematoda: <i>Anisakis</i> sp. (Whaleworm)	First intermediate host crustacean Euphasiid, second intermediate hosts fish that act as transport or paratentic hosts. Adult worms found in cetaceans.	Visceral cavity, musculature
<i>Pseudoterranova decipiens</i> (Sealworm)	First intermediate hosts crustacean mysids, second intermediate hosts fish that act as transport or paratentic hosts. Adult worms found in seals.	Visceral cavity, musculature
Digenea: <i>Cryptocotyle lingua</i> (Black Spot)	First intermediate host periwinkle, second intermediate host fish. Sea birds are definitive hosts.	Skin, subcutaneous tissue
Copepoda: <i>Caligus elongatus</i> (Sea Louse)	Two free-swimming naupliar, one infective copepodid, and four chalimus stages lead to the final adult stage on fish.	Body surface
<i>Lernaeocera branchialis</i>	Copepodid stage on flounder and lemon sole. Definitive host is cod, from which the female adult copepods disperse eggs.	Gills


Pathology

Most individual fish in wild populations are infested with parasites, but in the majority of cases, no significant harm appears to ensue. Parasites directly affect the health of the stock and spoil the appearance of the product, therefore reducing its market value.

Lernaeocera branchialis

The parasitic copepod (*Lernaeocera branchialis*), a parasite of the gills of cod, is probably the most serious pathogen of wild cod. Parasitized fish show markedly decreased condition and reduced deposition of fat in the skeletal muscle. Once settled on the gills of cod, adult female *L. branchialis* push their heads deeply into the hosts' tissues, damaging the heart and causing anemia.

Risks to farmed fish could occur with the placing of fish less than 1 year old at sites where the intermediate host (flat fish) is present in early summer/autumn, when the cod are particularly susceptible to infection.



Caligus elongatus, a sea louse. Photo courtesy of I. R. Bricknell.

***Cryptocotyle lingua* (black spot)**

Cryptocotyle lingua, or black spot, is a common parasite of cod that visit rocky shores during the summer. Juvenile cultured cod could be killed if free-swimming infective cercaria released from the intermediate periwinkle host penetrate the brains or hearts of the cod. The marketing of fish can also be hindered if large numbers of parasite-induced black spots cover the skins or surfaces of fish fillets.

Caligus elongatus

Sea lice (*Caligus elongates*) are a copepod crustacean parasite of many marine species including Atlantic cod. Through their attachment and feeding, copepodid and chalimus stages cause variable levels of localized damage. When present in high numbers, the parasites can cause significant pathology that results in mortality.

Ascaridoid Nematodes

Anisakis species and *Pseudoterranova decipiens*, both principal species involved in human infection by the family Anisakidae, have the potential to infect the flesh of cod. The parasites show limited infection within fish, but humans can become incidental hosts if fish are eaten raw or undercooked.

Control


The challenges presented to control parasites in Atlantic cod cultivation are multifaceted and must take into account a large number of variables. Transmission of parasites can occur directly, via vectors, or through environmental media such as water.

The main threat to farmed cod from parasites comes from those with direct or a single-host life cycle, such as *Caligus elongatus*, transmission of which is favored by the dense concentration of fish in intensive farming situations. In contrast, parasites with an intermediate host such as *Cryptocotyle lingua* undergo asexual reproduction within the intermediate host, the periwinkle. This increase in biotic potential can make control efforts more difficult.

The development, implementation, and ongoing enforcement of site-specific health management and biosecurity protocols are essential for the long-term financial and operational sustainability of cod farming. Careful site selection and improved pest management can minimize potential impacts from these pathogens.

Integrated control is an overall approach using strategically employed tactics based on knowledge of the life cycle and ecology of the parasite. It avoids overdependence on a single method and recognizes that no single method is likely to be 100 percent effective.

(Editor's Note: This article was originally published in the December 2005 print edition of the Global Aquaculture Advocate.)



Coiled *Anisakis* larvae encysted in the liver of a cod.

Authors



RACHEL KILBURN

Fisheries Research Services
FRS Marine Laboratory
P.O. Box 101, 375 Victoria Road
Aberdeen AB11 9DB
Scotland

kilburnr@marlab.ac.uk (<mailto:kilburnr@marlab.ac.uk>).



CAMPBELL PERT

Fisheries Research Services
FRS Marine Laboratory
P.O. Box 101, 375 Victoria Road
Aberdeen AB11 9DB
Scotland



DR. DAVID BRUNO

Fisheries Research Services
FRS Marine Laboratory
P.O. Box 101, 375 Victoria Road
Aberdeen AB11 9DB
Scotland



DR. ROB RAYNARD

Fisheries Research Services
FRS Marine Laboratory
P.O. Box 101, 375 Victoria Road
Aberdeen AB11 9DB
Scotland

Copyright © 2016–2020 Global Aquaculture Alliance

All rights reserved.