



[FEED SUSTAINABILITY \(/ADVOCATE/CATEGORY/FEED-SUSTAINABILITY\)](#)

Testing dietary potassium diformate in Pacific white shrimp

Monday, 31 July 2017

By Dr. Christian Lückstädt

Results show improved growth, survival, productivity in hatchery and grow-out conditions



Inclusion of potassium diformate in diets for postlarval and juvenile *L. vannamei* can improve survival, growth and productivity. Photo by Darryl Jory.

Intensive production of the Pacific or whiteleg shrimp (*Litopenaeus vannamei*) in Central America and Southeast Asia is estimated at over 3 million metric tons. Despite remarkable progress in shrimp nutrition and feed formulation during the past years, disease outbreaks in shrimp ponds can still lead to farming setbacks and, in some cases, to increased use of antibiotics.

Growing awareness from consumers and producers of aquacultured species is driving demand for responsible and sustainable aquaculture. Regulatory authorities in many seafood exporting countries now focus on the misuse of antibiotic growth promoters (AGP) in aquaculture, while public attention has shifted towards sustainable production methods. Alternative additives are being developed to replace AGPs.

Acidifiers are one of various alternatives spearheading environmental friendly and nutritive-sustainable aquaculture approaches. Currently, the most widely tested organic acid molecule in aquaculture is potassium diformate (KDF, Aquaform[®], ADDCON). It has been tested and used successfully in various cultured aquatic species, including salmon, trout, tilapia, Asian and European seabass, and pangasius. Its value to the shrimp production cycle has also been demonstrated in several field and research trials.

Potassium diformate is a double-salt formic acid molecule which decreases gastro-intestinal pH and thereby intensifies release of buffering fluids, containing enzymes, from the hepatopancreas. Formate also diffuses into pathogenic bacteria inside the digestive tract and acidifies their metabolism, leading to bacterial cell death. Furthermore, beneficial bacteria (*Lactobacilli*, *Bifidobacteria*) are supported (*eubiosis*), which may lead to improved gut health, resulting in stronger condition of the shrimp.

One of the most crucial periods in the life cycle of shrimp is the post-larval stage, when shrimp feeding changes from algae and brine shrimp nauplii, to commercially formulated larval diets. Shrimp survival rates during this period are critical to later productivity, but various pathogenic bacteria can significantly increase larval mortality in shrimp hatcheries.

Dietary potassium diformate and *L. vannamei* postlarvae and juveniles

Dr. He and colleagues in China (2006) carried out an experiment to test the use of potassium diformate on *L. vannamei* postlarvae (average body weight 57 mg) through a formulated diet containing either zero or 0.8 percent potassium diformate (KDF) fed for a period of 40 days. Shrimp larvae fed the diet with KDF had a significantly improved performance in terms of growth, feed conversion and survival rates (Table 1).

Lückstädt, KDF, Table 1

Parameter	Control	0.8 percent KDF	Difference (%)
Initial BW (mg)	57	57	–
Final BW (mg)	256±34	309±35	+21
WG [mg, 40 d]	199	252	+27
FCR	3.73±0.6	2.49±0.3	-33
Survival (%)	92.2±1.6	100	+8
Prod. Index*	0.49	1.01	+106

Growth, feed conversion and survival of *Litopenaeus vannamei* PL under aquaria conditions after 40 days.

*Productivity Index (weight gain [g] × survival [percent] / (10 × FCR)).

Shrimp larvae fed the diet with KDF inclusion showed a more efficient growth and significantly improved feed utilization, as well as lower mortality ($P < 0.05$), compared to control animals, resulting in a markedly increased productivity index, which is a formula including the three most important production figures in shrimp production: weight gain of shrimp, feed efficiency and survival rate. Optimized nutrition in the early stages of the shrimp production cycle often results in an overall improved shrimp productivity, which was also demonstrated in another experiment reported by Jintasataporn et al. (2011).

We carried out another experiment in aquaria to simulate intensive growout and investigate the growth performance of juvenile *L. vannamei* shrimp fed with low levels of dietary KDF vs. a control diet. For the 10-week experiment, 30 aquaria (120-L each) and connected to a flow-through system, were filled with 20 ppt seawater and 28.0 ± 2.0 degrees-C and > 6.0 ppm of dissolved oxygen. Each aquarium was stocked with 18 shrimp each, with an average body weight of 2.4 ± 0.1 g ($n=540$). Shrimp were fed to satiation three times a day, with a commercial diet containing 32 percent crude protein. The control diet (treatment A) did not contain KDF, and 0.2 percent and 0.5 percent KDF were added to diets in treatments B and C, respectively.

Results showed higher individual body mass, daily weight gain and specific growth rates for shrimp fed on the diets that included KDF (treatments B+C). Those KDF-fed shrimp reached growth parameters compared to the control group (treatment A) of 11.8 g vs. 11.0 g final weight, 0.13 g/ind/day vs. 0.12 g/ind/day and an SGR of 2.26 percent/day vs. 2.16 percent/day, respectively (Table 2).

Similarly, the survival rates of shrimp fed diets with KDF were 80.6 vs. 76.1 percent in the control group, and feed conversion ratios (FCR) were 1.37 vs. 1.47. Thus, the growth performance of Pacific white shrimp fed diets with 0.2 and 0.5 percent KDF inclusions resulted in significantly ($P < 0.05$) increased body weight by 7.2 and 7.4 percent, and higher average daily weight gains of 9.26 and 9.17 percent ($P = 0.06$). Similarly, FCR values were improved by 7.1 and 7.0 percent ($P = 0.07$) compared to the control group.

Lückstädt, KDF, Table 2

Parameter	Neg. control	0.2% KDF	0.5% KDF
Initial BM (g/ind)	2.4 ± 0.1	2.4 ± 0.1	2.5 ± 0.1
Final BM (g/ind)	$11.0b \pm 0.8$	$11.8a \pm 1.3$	$11.8a \pm 0.7$
Weight gain (g)	8.6 ± 0.9	9.4 ± 1.3	9.4 ± 0.7
SGR (%)	2.2 ± 0.1	2.3 ± 0.2	2.3 ± 0.1
Survival (%)	76.1 ± 7.0	76.1 ± 4.6	80.6 ± 13.4
FCR	1.47 ± 0.1	1.37 ± 0.1	1.37 ± 0.1
Productivity Index (PI)	$45.0b \pm 8.5$	$53.8ab \pm 14.5$	$55.9a \pm 14.0$

Performance of whiteleg shrimp fed with or without dietary KDF. SGR = Specific Growth Rate; FCR = Feed Conversion Ratio; PI = Weight gain (g) x Survival (percent) / (FCR x 10). Means with different superscripts within rows are significantly different ($P < 0.05$).

If the data are analyzed for overall productivity, using the Productivity Index, it shows that the inclusion of dietary KDF resulted in significant improvements of the PI vs. the negative control by more than 19 percent or 24 percent, respectively. We conclude that using our dietary potassium diformate is a promising alternative in modern shrimp nutrition that contributes to an economically and ecologically sustainable growout operation.

Other experiments carried out under controlled laboratory conditions, like the one discussed above, showed unusually high survival rates (76 to 81 percent), and thus were not mimicking bacterial situations on commercial farms closely enough. Survival rates that do not reflect those found in commercial farms often provide an unrealistic picture of the additive's benefit. Under commercial farm conditions, shrimp can be exposed to a number of different challenges, including bacterial pathogens.

Vibrio challenge trial

Therefore, a subsequent trial was carried out to challenge juvenile *L. vannamei* shrimp with the bioluminescent, Gram-negative bacterium *V. harveyi*, which regularly causes increased mortality in shrimp culture. The trial consisted of a negative control compared against two treatment groups (0.2 percent and 0.5 percent dietary KDF, at dosages similar to the ones used in the trial reported above). A total of 90 shrimp (30 shrimp per group), with a mean body weight of 11.0 ± 0.8 g, were used. The trial used the same protocols as described above, but with the addition of the pathogenic *V. harveyi* to the water at the beginning of the 10-day trial at a concentration of 5×10^6 CFU/mL.



Dietary inclusion of potassium diformate improved the survival of juvenile *L. vannamei* challenged with *Vibrio harveyi*. Photo by Darryl Jory.

At the end of this challenge trial, the mortality in the non-treated shrimp was significantly higher ($P < 0.01$; 76.6 ± 5.8 percent) compared to shrimp fed with the diets with KDF at both inclusion levels (50.0 ± 10.0 percent for both 0.2 percent and 0.5 percent KDF), as shown in Fig. 1. The effect of the acidifier was clear from the first day of the trial, but the difference between treatment and control became more pronounced from day 4 onwards and remained significantly different till the end of the trial. Both dietary KDF dosages reduced the mortality in the challenged shrimp to the same extent by day 10.




Fig. 1. Mortality of *L. vannamei* shrimp fed potassium diformate (KDF, AQUAFORM®, ADDCON) under challenge with the pathogenic bacterium *Vibrio harveyi*.

Perspectives

We conclude from our research that our dietary potassium diformate can reduce mortality in Pacific white shrimp caused by the Gram-negative, pathogenic bacterium *V. harveyi*. It may be expected that similar outcomes may result with other Gram-negative bacterial pathogens in commercial shrimp farming operations.

Considering our results on growth performance and survival rates, we believe that potassium diformate is a promising additive for economic and sustainable shrimp production, and should be considered in compound feeds for commercial, outdoor shrimp farming operations.

References available from author.

Author



DR. CHRISTIAN LÜCKSTÄDT

ADDCON GmbH

Parsevalstrasse 6

06749 Bitterfeld-Wolfen

christian.lueckstaedt@addcon.com (<mailto:christian.lueckstaedt@addcon.com>).

Copyright © 2016–2018
Global Aquaculture Alliance